

Policy Name	Clinical Policy – Glaucoma Surgery
Policy Number	1327.00
Department	Clinical Product & Development
Subcategory	Medical Management
Original Approval Date	05/01/2018
Current MPC/CMO Approval Date	04/03/2024
Current Effective Date	07/01/2024

Company Entities Supported (Select All that Apply)

- X Superior Vision Benefit Management
- X Superior Vision Services
- \overline{X} Superior Vision of New Jersey, Inc.
- X Block Vision of Texas, Inc. d/b/a Superior Vision of Texas
- X Davis Vision

(Collectively referred to as 'Versant Health' or 'the Company'

Acronyms and Definitions		
IOP	Intraocular pressure	
MIGS	Minimally Invasive Glaucoma Surgery	
Target Pressure	The upper limit of a range of pressures in which visual field loss is unlikely to risk a patient's vision, health or quality of life.	

PURPOSE

To provide the medical necessity criteria to support the indication(s) for glaucoma surgery and to render medical necessity determinations. Applicable procedure codes related to glaucoma surgery are also defined.

POLICY

A. BACKGROUND

Glaucoma is an optic neuropathy that can lead to progressive visual field loss and significant impairment of both vision and health related quality of life. Reducing intraocular pressure is



associated with a decreased risk of developing or progressing optic nerve damage. There is no absolute number or range of intraocular pressure (IOP) that defines glaucoma.1

Glaucoma surgical procedures are intended to lower IOP longer term, thus reducing the risk of permanent optic nerve damage. IOP can be lowered in a variety of ways. One method is by enhancing the outflow of aqueous from the anterior chamber through various methods. Some types of procedures open, enhance or stent the eye's natural drainage system usually by reducing obstruction at the level of the trabecular meshwork. A second method is to create an alternative route for aqueous drainage² through either a microshunt, fistulization of the sclera, or implantation of a glaucoma drainage implant. A third method is to reduce the amount of aqueous humor production through partial destruction of the ciliary body.3

B. Medically Necessary

- 1. Minimally Invasive Glaucoma Surgery
 - a. Select aqueous drainage assist devices with concurrent cataract extraction and implantation of an intraocular lens (66989/66991): Please refer to Versant Health Clinical Policy 1301 Cataract Surgery for the medical necessity criteria guidelines.
 - b. Excisional (Trabeculotomy, Goniotomy, GATT 65820)
 - i. Initial treatment for congenital or juvenile glaucoma. In this situation, does not require a trial of topical therapy, or;
 - ii. Mild to moderate open angle glaucoma; and,
 - iii. When a trial of medical therapy (medication and laser therapy) has been unable to control intraocular pressure and reduce the risk for further optic nerve damage due to lack of efficacy, compliance, cognitive impairment, or coexisting ocular surface disease.
 - c. Transluminal dilation of aqueous outflow canal (66174, 66175)
 - Diagnosis of mild to moderate open angle glaucoma confirmed by gonioscopy; and.
 - ii. When a trial of medical therapy (medication and laser therapy) has been unable to control intraocular pressure and reduce the risk for further optic nerve damage due to lack of efficacy, compliance, cognitive impairment, or coexisting ocular surface disease.
 - d. Ab interno implant (Xen 0449T/0450T)
 - i. Diagnosis of mild to moderate open angle glaucoma confirmed by gonioscopy;
 - ii. When a trial of medical therapy (medication and laser therapy) has been unable to control intraocular pressure and reduce the risk for further optic nerve damage

¹ Sommer, 2011

² See Clinical Policy 1308 Laser Trabeculoplasty; and Clinical Policy 1322 Laser Peripheral Iridotomy.

³ See Clinical Policy 1300 Cataract Surgery for procedures done in conjunction with cataract surgery



due to lack of efficacy, compliance, cognitive impairment, or coexisting ocular surface disease.

- Trabeculectomy and related fistulization procedures, with or without stents or drainage device (Ab externo implant: ExPRESS, Ab externo XEN [66183]; Trabeculectomy [66170,66172]; Tube Shunt: Ahmed, Baerveldt, Molteno [66179/66180]
 - a. Diagnosis of glaucoma; and,
 - b. Maximum medical therapy (maximum topical therapy and if appropriate laser therapy) has been unable to control intraocular pressure and reduce the risk for further optic nerve damage due to lack of efficacy, compliance, or cognitive impairment.

C. Not Medically Necessary

- 1. Glaucoma Surgery is not medically necessary as follows:
 - a. In the absence of glaucoma; or,
 - b. When tolerated medication provides adequate control of intraocular pressure; or,
 - c. When evidence of visual field loss or nerve fiber layer deficit is not documented; or,
 - d. For ocular hypertension or glaucoma suspect; or,
 - e. For off-label use of an aqueous drainage device or glaucoma drainage device; or,
 - f. When there is no risk for further optic nerve damage.
- 2. Glaucoma drainage devices and aqueous drainage assist devices not explicitly approved by the FDA are considered experimental and/or investigational and may not be considered medically necessary.

D. Documentation

Medical necessity must be supported by adequate and complete documentation in the patient's medical record that describes the procedure and the medical rationale. Documentation requires at a minimum all the following items. All items must be available upon request. For any retrospective review, a full operative report is needed.

Every page of the record must be legible and include appropriate patient identification information (e.g., complete name, date(s) of service). Services provided/ordered must be authenticated by the physician. The method used shall be handwritten or electronic signature. Stamped signatures are not acceptable.

- Eye exam with description of medical justification for the proposed glaucoma surgical procedure and absence of contraindications for the surgery. This must include gonioscopy and thorough evaluation of the optic nerve to validate structural glaucomatous changes; and,
- 2. Documentation of glaucoma findings and inability to reduce risk for optic nerve damage with medical therapy due to noncompliance with medical therapy such as cost, memory problems, difficulty with instillation, adherence, or intolerance to the medication; and,



- 3. Documentation of ongoing risk of optic nerve damage despite topical therapies and laser trabeculoplasty, when considering trabeculectomy and related external fistulization procedures with or without stents or other drainage devices; and,
- 4. Allied diagnostic testing with physician's order, medical rationale, findings, interpretation and report demonstrating reproducible visual field loss with threshold perimetry, and optic disc changes on OCT and/or fundus photography. The perimetry requirement is waived for patients who cannot reliably perform this test (e.g., nystagmus); and,
- 5. Documentation of previously failed glaucoma surgery when applicable; and,
- 6. A statement from the surgeon anticipating the procedure(s) will achieve reduction of risk to the optic nerve, and that this reduction will be adequate to control the glaucoma; and,
- 7. An explanatory statement from the surgeon when more than one aqueous drainage assist device, or glaucoma drainage device, will be implanted in the same eye during the same operative session; and,
- 8. For retrospective review, detailed operative report that incorporates indications, procedure description, make, model, serial number(s) of implant(s) that are FDA approved.

E. PROCEDURAL DETAIL

CPT Codes		
0449T	Insertion of aqueous drainage device, without extraocular reservoir, internal approach, into the subconjunctival space; initial device	
0450T	Insertion of aqueous drainage device, without extraocular reservoir, internal approach, into the subconjunctival space; each additional device (List separately in addition to code for primary procedure) Add on code for 0449T	
0474T	Insertion of anterior segment aqueous drainage device, with creation of intraocular reservoir, internal approach, into the supraciliary space	
0671T	Insertion of anterior segment aqueous drainage device into the trabecular meshwork, without external reservoir, and without concomitant cataract removal, one or more	
65820	Goniotomy	
66150	Fistulization of sclera for glaucoma; trephination with iridectomy	
66155	Fistulization of sclera for glaucoma; thermocauterization with iridectomy	
66160	Fistulization of sclera for glaucoma; sclerectomy with punch or scissors, with iridectomy	
66170	Fistulization of sclera for glaucoma; trabeculectomy ab externo in absence of previous surgery	
66172	Fistulization of sclera for glaucoma; trabeculectomy ab externo with scarring from previous ocular surgery or trauma (includes injection of antifibrotic agents)	
66174	Transluminal dilation of aqueous outflow canal; without retention of device or stent	
66175	Transluminal dilation of aqueous outflow canal; with retention of device or stent	
66179	Aqueous shunt to extraocular equatorial plate reservoir; external approach; without graft	



66180	Aqueous shunt to extraocular equatorial plate reservoir; external approach; with graft.
66183	Insertion of anterior segment aqueous drainage device, without extraocular reservoir, external approach (also known as express shunt).
66184	Revision of aqueous shunt to extraocular equatorial plate reservoir; without graft
66185	Revision of aqueous shunt to extraocular equatorial plate reservoir; with graft
66250	Revision or repair of operative wound of anterior segment, any type, early or late, major or minor procedure
66989	Extracapsular cataract removal with insertion of intraocular lens prosthesis (1-stage procedure), manual or mechanical technique (e.g., irrigation and aspiration or phacoemulsification), complex, requiring devices or techniques not generally used in routine cataract surgery (e.g., iris expansion device, suture support for intraocular lens, or primary posterior capsulorrhexis) or performed on patients in the amblyogenic developmental stage; with insertion of intraocular (e.g., trabecular meshwork, supraciliary, suprachoroidal) anterior segment aqueous drainage device, without extraocular reservoir, internal approach, one or more.
66991	Extracapsular cataract removal with insertion of intraocular lens prosthesis (1 stage procedure), manual or mechanical technique with insertion of intraocular, anterior segment aqueous drainage device, without extraocular reservoir, internal approach, one or more.
Required	Modifiers
Anatomic al Modifiers	RT – Right side LT – Left side 50 – Bilateral
Invalid Mo	odifiers
EM Modifiers	24, 25, 57, 95
Diagnosti c Modifiers	26, TC
Other Modifiers	E1, E2, E3, E4, JW, KX

DISCLAIMER and COPYRIGHTS

This clinical policy is provided for information purposes only and does not constitute medical advice. Versant Health, Inc., and its affiliates (the "Company") do not provide health care services and cannot guarantee any results or outcomes. Treating doctors are solely responsible for determining what services or treatments to provide to their patients. Patients (members) should always consult their doctor before making any decisions about medical care.

Subject to applicable law, compliance with this clinical policy is not a guarantee of coverage or payment. Coverage is based on the terms of an individual's particular benefit plan document,



which may not cover the service(s) or procedure(s) addressed in this clinical policy. The terms of the individual's specific benefit plan are always determinative.

Every effort has been made to ensure that the information in this clinical policy is accurate and complete, however the Company does not guarantee that there are no errors in this policy or that the display of this file on a website is without error. The company and its employees are not liable for any errors, omissions, or other inaccuracies in the information, product, or processes disclosed herein. Neither the company nor the employees represent that the use of such information, products, or processes will not infringe on privately owned rights. In no event shall the Company be liable for direct, indirect, special, incidental, or consequential damages arising out of the use of such information, product, or process.

COMPANY'S COPYRIGHT STATEMENT Except for any copyrights described below, this clinical policy is confidential and proprietary, and no part of this clinical policy may be copied, distributed or used without Versant Health, or its applicable affiliates' express prior written approval.

AMA COPYRIGHT STATEMENT CPT© is the 2002-2024 copyright of the American Medical Association. All Rights Reserved. CPT™ is a registered trademark of the American Medical Association. Applicable FARS/DFARS apply to government use. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

RELATED POLICIES AND PROCEDURES		
1300	Cataract Surgery	
1308	Laser Trabeculoplasty	
1322	Laser Peripheral Iridotomy	

DOCUMENT HISTORY			
Approval Date	Revisions	Effective Date	
05/01/2018	Initial Policy	05/01/2018	
03/13/2019	Annual review; no criteria change.	03/13/2019	
10/18/2019	Correction of code 66160.	11/01/2019	
08/19/2020	Annual review; no criteria change.	01/01/2021	
07/07/2021	Annual review; added background information on IOP measurement; removed exclusion from surgery for closed angle glaucoma and excessive synechiae; removed inclusion of borderline glaucoma; added requirement for presurgical eye exam to include gonioscopy.	01/01/2022	



10/06/2021	Revised criteria for surgery to not require visual field loss due to glaucoma.	01/01/2022
01/05/2022	Annual review; revised criteria for MIGS to match criteria in 1300 for same procedure codes. Remove two CMS deleted codes; add three CMS released codes.	04/01/2022
01/04/2023	Annual review; revised language related to achieving or requiring measures of target intraocular pressure to now describe risk/status of optic nerve damage as primary measurement for surgery; added goniotomy as medically necessary test for three indications; removed requirement to submit surgical clearance documentation.	07/01/2023
09/20/2023	Administrative review for CMS 2024 final rule Medicare Part C equity: no changes.	n/a
04/03/2024	Removed MIGS combination cataract surgery criteria which is retained in 1300 Cataract Surgery policy. Add indication of laser trabeculoplasty failure for trabeculotomy/goniotomy procedures; added indication of risk to the optic nerve with further laser therapy for procedures of trabeculectomy and related fistulization	07/01/2024

REFERENCES AND SOURCES

- 1. Abdullah S, Jasek MC, Radcliffe NM, et al. A novel dual blade device for goniotomy: initial clinical experience. Invest Ophthalmol Vis Sci. 2016; 57: 6522.
- Ahmed IIK, Fea A, Au L, et.al. A Prospective Randomized Trial Comparing Hydrus and iStent Microinvasive Glaucoma Surgery Implants for Standalone Treatment of Open-Angle Glaucoma: The COMPARE Study. Ophthalmology. 2020 Jan;127(1):52-61. doi: 10.1016/j.ophtha.2019.04.034. Epub 2019 Apr 26. PMID: 31034856.
- 3. Andrew NH, Akkach S, Casson RJ. A review of aqueous outflow resistance and its relevance to microinvasive glaucoma surgery. Survey Ophthalmol. 2020;65(1):18-31. doi: 10.1016/j.survophthal.2019.08.002.
- 4. Arona K.S., Robin A.L., Corcoran K.J., et. al., Use of Various Glaucoma Surgeries and Procedures in Medicare Beneficiaries from 1994 to 2012,
- 5. Aref AA, Gedde SJ, Budenz DL. Glaucoma Drainage Implant Surgery. Dev Ophthalmol. 2017; 59:43-52. doi: 10.1159/000458485. Epub 2017 Apr 25. PMID: 28442686.
- 6. Berdahl J, Voskanyan L, Myers JS, et al. Implantation of two second-generation trabecular micro-bypass stents and topical travoprost in open-angle glaucoma not controlled on two preoperative medications: 18- month follow-up. *Clinical & Experimental Ophthalmology.* 2017;45(8):797-802.
- 7. Brusini P., "Canaloplasty in Open-Angle Glaucoma Surgery: A Four-Year Follow-Up," The Scientific World Journal, vol. 2014, Article ID 469609, 7 pages, https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3915493/.
- 8. Brusini, P, Caramello, G, Benedetti, S, Tosoni, C, "Canaloplasty in open angle glaucoma. Mid-term results from a multicenter study," Journal of Glaucoma 2016 May; 25(5):403-7.
- 9. Budenz DL, Gedde SJ. New options for combined cataract and glaucoma surgery. Curr Opin Ophthalmol. 2014 Mar;25(2):141-7. doi: 10.1097/ICU.000000000000027. PMID:



- 24389806.
- 10. Bussel II, Kaplowitz K, Schuman J. et.al. Study Group (2015) Outcomes of ab interno trabeculectomy with the trabectome after failed trabeculectomy. Br J Ophthalmol. 2015. 99: 258-262.
- 11. Chaudhary A, Salinas L, Guidotti J, et.al. XEN Gel Implant: a new surgical approach in glaucoma. Expert Rev Med Devices. 2018 Jan;15(1):47-59. doi: 10.1080/17434440.2018.1419060. Epub 2017 Dec 26. PMID: 29258404.
- 12. Chen TC, Chen PP, Francis BA, Junk AK, Smith SD, Singh K, Lin SC. Pediatric glaucoma surgery: a report by the American Academy of Ophthalmology. Ophthalmology. 2014 Nov;121(11):2107-15. doi: 10.1016/j.ophtha.2014.05.010. Epub 2014 Jul 24. PMID: 25066765.
- 13. Conlon R, Saheb H, Ahmed II. Glaucoma treatment trends: a review. Can J Ophthalmol. 2017 Feb;52(1):114-124. doi: 10.1016/j.jcjo.2016.07.013. Epub 2016 Nov 17. PMID: 28237137.
- 14. Craven ER, Katz LJ, Wells JM, et al. iStent Study Group. Cataract surgery with trabecular micro-bypass stent implantation in patients with mild-to-moderate open-angle glaucoma and cataract: Two-year follow up. *J Cataract Refract Surg.* 2012 Aug; 38(8):1339-45.
- 15. De Gregorio A, Pedrotti E, Russo L, Morselli S. Minimally invasive combined glaucoma, and cataract surgery: clinical results of the smallest ab interno gel stent. *Int Ophthalmol.* 2017:1-6. doi: 10.1007/s10792-017-0571.
- 16. DeJong L, Lafuma A, Aguadé AS, et.al. Five-year extension of a clinical trial comparing the EX-PRESS glaucoma filtration device and trabeculectomy in primary open-angle glaucoma. Clin Ophthalmol. 2011;5: 527-533.
- 17. Do AT, Parikh H, Panarelli JF. Subconjunctival microinvasive glaucoma surgeries: an update on the Xen gel stent and the PreserFlo MicroShunt. Curr Opin Ophthalmol. 2020 Mar;31(2):132-138. doi: 10.1097/ICU.000000000000643. PMID: 31922979.
- 18. Donnenfeld ED, Solomon KD, Voskanyan L, et al. A prospective 3-year follow-up trial of implantation of two trabecular micro bypass stents in open-angle glaucoma. *Clinical Ophthalmology*. 2015; 9:2057-2065.
- 19. Epstein RS, Scott AT, Capitena Young CE, et.al. Optimum time for angle visualization during ab interno glaucoma surgery: Before or after phacoemulsification. J Cataract Refract Surg. 2019 May;45(5):615-619. doi: 10.1016/j.jcrs.2019.01.012. PMID: 31030776.
- 20. Fellman R.L., Mattox C., Singh K., et. al., American Glaucoma Society Position Paper: Microinvasive Glaucoma Surgery, Ophthal Glaucoma Jan-Feb 2020; 3(1):1-6
- 21. Ferguson TJ, Berdahl JP, Schweitzer JA, Sudhagoni R. Evaluation of a trabecular microbypass stent in pseudo phakic patients with open-angle glaucoma. *Journal of glaucoma*. 2016;25(11):896-900.
- 22. Francis BA, Hong B, Winarko J, Kawji S, et. al., Vision loss and recovery after trabeculectomy: risk and associated risk factors. Arch Ophthalmology 2011: 129(8):1011-1017.
- 23. Galal A, Bilgic A, Eltanamly R, et.al. XEN Glaucoma Implant with Mitomycin C 1-Year Follow-Up: Result and Complications. *J Ophthalmol.* 2017; 2017:5457246.
- Geffen N, Assia EI, Melamed S. Laser-Assisted Techniques for Penetrating and Nonpenetrating Glaucoma Surgery. Dev Ophthalmol. 2017; 59:100-112. doi: 10.1159/000458490. Epub 2017 Apr 25. PMID: 28442691.
- 25. Gillmann K, Mansouri K. Minimally Invasive Glaucoma Surgery: Where Is the Evidence? Asia Pac J Ophthalmol (Phila). 2020 May-Jun;9(3):203-214. doi: 10.1097/APO.000000000000294. PMID: 32501895; PMCID: PMC7299223.
- 26. Greenwood MD, Seibold L, et al. Goniotomy with a single-use blade: Short-term results. Journal of Cataract & Refractive Surgery. 2017: 43 (9) 1197-1201.



- 27. Grover DS, Flynn WJ, Bashford KP, et al. Performance and Safety of a New Ab Interno Gelatin Stent in Refractory Glaucoma at 12 Months. *Am J Ophthalmol*. 183:25-36. 2017.
- 28. Grover DS, Godfrey DG, Smith O, et.al. Gonioscopy-assisted transluminal trabeculotomy, ab interno trabeculotomy: technique report and preliminary results. Ophthalmology. 2014. 121: 855-861.
- 29. Grover DS, Godfrey DG, Smith O, et.al. Outcomes of Gonioscopy-assisted Transluminal Trabeculotomy (GATT) in Eyes with Prior Incisional Glaucoma Surgery. J Glaucoma. 2017. 26: 41-45.
- 30. Hohberger B, Welge-LüBen UC, Lämmer R. ICE-Syndrome: A case report of implantation of a microbypass Xen gel stent after DMEK transplantation. *J Glaucoma*. 2017; 26(2): e103–e104.
- 31. Hu R, Racette L, Chen KS, et. al., Functional Assessment of glaucoma: uncovering progression, Survey of Ophthalmology 639-661, August 2020.
- 32. Jabłońska J, Lewczuk K, Konopińska J, Mariak Z, Rękas M. Microinvasive glaucoma surgery: a review and classification of implant-dependent procedures and techniques. Acta Ophthalmol. 2022 Mar;100(2):e327-e338. doi: 10.1111/aos.14906. Epub 2021 May 14. PMID: 33988310; PMCID: PMC9291507.
- 33. Junoy Montolio FG, Müskens RPHM, Jansonius NM. Influence of glaucoma surgery on visual function: a clinical cohort study and meta-analysis. Acta Ophthalmol. 2019 Mar;97(2):193-199. doi: 10.1111/aos.13920. Epub 2018 Oct 4. PMID: 30288923; PMCID: PMC6586003.
- 34. Kang JM, Tanna AP. Glaucoma. Med Clin North Am. 2021 May;105(3):493-510. doi: 10.1016/j.mcna.2021.01.004. Epub 2021 Apr 2. PMID: 33926643.
- 35. Katz LJ, Erb C, Carceller GA, et al. Prospective, randomized study of one, two, or three trabecular bypass stents in open-angle glaucoma subjects on topical hypotensive medication. *Clinical Ophthalmology*. 2015; 9:2313-2320.
- 36. Khaimi, MA. Ab Interno Canaloplasty. Glaucoma Today. http://glaucomatoday.com/pdfs/gt1115_surgpearls.pdf November/December 2015.
- 37. Kornmann HL, Gedde SJ. Glaucoma management after corneal transplantation surgeries. Curr Opin Ophthalmol. 2016 Mar;27(2):132-9. doi: 10.1097/ICU.0000000000000237. PMID: 26595847; PMCID: PMC5584078.
- 38. Lee JH, Amoozgar B, Han Y (2017) Minimally Invasive Modalities for Treatment of Glaucoma: An Update. J Clin Exp Ophthalmol. 2017. 8: 666.
- 39. Lewis RA, von Wolff K, Tetz M, et al. Canaloplasty: three-year results of circumferential vasodilation and tensioning of Schlemm's canal using a micro catheter to treat open-angle glaucoma. J Cataract Refract Surg. 2011; 37:682-690.
- 40. Lin MK, Ge J. How to make the intraocular pressure after glaucoma filtering surgery more controllable]. 2020 Jan 11;56(1):13-16. Chinese. doi: 10.3760/cma.j.issn.0412-4081.2020.01.005. PMID: 31937058.
- 41. Pérez-Torregrosa VT, Olate-Pérez Á, Cerdà-Ibáñez M, et al. Combined phacoemulsification, and Xen 45 surgery from a temporal approach and 2 incisions. *Arch Soc Esp Oftalmol.* 2016; 91(9):415-421.
- 42. Pinto FN, Abegã PL, Marques-Neves C. Xen gel stent internal ostium occlusion: ab-interno revision. *J Glaucoma*. 2017; 26(4): e150–e152.
- 43. Roelofs K, Arora S, Dorey MW. Implantation of 2 trabecular micro bypass stents in a patient with primary open-angle glaucoma refractory to previous glaucoma-filtering surgeries. *Journal of Cataract & Refractive Surgery.* 2014; 40(8):1322-1324.
- 44. Hu R, Racette L, Chen KS, Johnson CA. Functional assessment of glaucoma: Uncovering progression. Surv Ophthalmol. 2020;65(6):639-661. doi: 10.1016/j.survophthal.2020.04.004.



- 45. Samuelson TW, Katz LJ, Wells JM, et al. US iStent Study Group. Randomized Evaluation of the Trabecular Micro-Bypass Stent with Phacoemulsification in Patients with Glaucoma and Cataract. Ophthalmology 2011.
- 46. Schlenker MB, Gulamhusein H, Conrad-Hengerer I, et al. Efficacy, Safety, and Risk Factors for Failure of Standalone Ab Interno Gelatin Micro Stent Implantation versus Standalone Trabeculectomy. *Ophthalmology*. 2017; 124(11):1579-1588.
- 47. Somer A. Ocular Hypertension and Normal Tension Glaucoma, Arch of Ophthalmology;129,6, June 2011, p. 785-6.
- 48. Yang X, Zhao Y, Zhong Y, et.al. The efficacy of XEN gel stent implantation in glaucoma: a systematic review and meta-analysis. BMC Ophthalmol. 2022 Jul 15;22(1):305. doi: 10.1186/s12886-022-02502-y. PMID: 35836197; PMCID: PMC9284889.

SOURCES

- 1. American Academy of Ophthalmology, Primary Open Angle Glaucoma PPP 2020. https://www.aao.org/education/preferred-practice-pattern/primary-open-angle-glaucoma-ppp. Accessed 10/2023.
- 2. American Academy of Ophthalmology, Primary Open Angle Glaucoma Suspect PPP 2020. https://www.aao.org/education/preferred-practice-pattern/primary-open-angle-glaucoma-suspect-ppp. Accessed 10/2023.
- 3. 5. American Academy of Ophthalmology, Glaucoma Summary Benchmarks 2022. https://www.aao.org/education/summary-benchmark-detail/glaucoma-summary-benchmarks-2020. Accessed 10/2023.
- 4. American Academy of Ophthalmic Executives® Fact Sheet: Goniotomy. Jan 2023. https://www.aao.org/Assets/c1c5ad6a-f611-4c41-988c-991514f68602/637896975656770000/goniotomy-fs-pdf?inline=1. Accessed 11/2023.
- American Academy of Ophthalmic Executives® Fact Sheet: Coding for iStent, iStent inject, iStent infiniteand Hydrus/ Aug. 2023. https://www.aao.org/Assets/edd35b50-5596-49b2-8709-22316ead6c41/637975734180730000/istent-istent-inject-hydrus-fs-pdf?inline=1. Accessed 11/2023.
- American Academy of Ophthalmic Executives® 0449T XEN® Gel Stent. June 2023. https://www.aao.org/Assets/bd71150c-c9ee-4622-be06-78eda413afa7/637886642913100000/0449t-xen-gel-stent-fs-pdf?inline=1. Accessed 11/2023.
- CMS Micro Invasive Glaucoma Surgery (MIGS) DL37578 PROPOSED, July 2023.
 https://www.cms.gov/medicare-coverage-database/view/lcd.aspx?lcdid=39581&ver=6&keyword=MIGS&keywordType=starts&areald=all&docType=NCA,CAL,NCD,MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=1. Accessed 10/2023.