

Policy Name	Clinical Policy – Optical Coherence Tomography (OCT)
Policy Number	1304.00
Department	Clinical Product & Development
Subcategory	Medical Management
Original Approval Date	07/10/2017
Current MPC/CCO Approval Date	04/03/2024
Current Effective Date	06/01/2024

Company Entities Supported (Select All that Apply) <input checked="" type="checkbox"/> Superior Vision Benefit Management <input checked="" type="checkbox"/> Superior Vision Services <input checked="" type="checkbox"/> Superior Vision of New Jersey, Inc. <input checked="" type="checkbox"/> Block Vision of Texas, Inc. d/b/a Superior Vision of Texas <input checked="" type="checkbox"/> Davis Vision (Collectively referred to as 'Versant Health' or 'the Company')

ACRONYMS	
OCT	Optical Coherence Tomography
OCT-A	Optical Coherence Tomography Angiography

PURPOSE

To provide the clinical criteria to support the indication(s) for Optical Coherence Tomography (OCT). Applicable procedure codes are also defined.

POLICY

A. BACKGROUND

Optical coherence tomography is performed on the anterior or the posterior segment of the eye to identify and quantify structural changes associated with ocular disease, which may not be visible with other forms of ophthalmoscopy.

B. Medically Necessary

OCT may be medically necessary when the information garnered from an eye exam,

including routine ophthalmoscopy, is insufficient to assess the patient's disease.

1. Anterior OCT may be medically necessary to evaluate narrow angles and disorders of the cornea, iris and ciliary body. It may be necessary in measurements of abnormally thin corneas, visualization of malpositioned IOLs (intraocular lens) or other implants in the anterior segment, assessment of lesions on the anterior segment ocular structures, and measurement of a narrow anterior chamber.
2. Posterior OCT may be medically necessary to evaluate conditions of glaucoma, macular traction, diabetic retinopathy, macular hole, macular edema, age-related macular degeneration, and other retinal and optic nerve abnormalities
3. Posterior OCT may be medically necessary as a baseline study prior to initiation of chloroquine, hydroxychloroquine, or vigabatrin therapy. Patients should be monitored via OCT-P for drug-induced retinopathy within the first year of treatment and annually thereafter starting at year five of use. For high risk patients, annual testing may begin immediately without a 5-year delay¹.

C. Not Medically Necessary

OCT may not be medically necessary for the following:

1. To evaluate an eye without signs, symptoms, serious ophthalmic disease, ocular abnormalities, or contributory medical history such as high-risk long-term drug therapy.
2. When performed during the global surgery period of an ophthalmic surgical procedure to verify the expected outcome. For example, OCT after macular pucker surgery, may not be medically necessary to verify success of the procedure.
3. Without a documented medical rationale in the medical record
4. To confirm a diagnosis that has already been determined
5. When other related ophthalmological tests (e.g., fundus photography, angiography, ultrasound, etc.) have been performed, OCT may not be medically necessary unless it provides additive and not duplicative information.
6. OCT of the optic disc or retina at the same time as a fundus photograph is not medically necessary unless the medical record shows how the test results for each test were required for the patient's plan of care in an additive and non duplicative manner.
7. When posterior OCT and extended ophthalmoscopy (CPT 92201, 92202) are performed concurrently, the retinal drawing must identify severe posterior segment disease not identifiable in the image. Otherwise, the procedures are considered duplicative, and only one of the procedures may be considered medically necessary.

¹ AAO Drug Induced Maculopathy, 2023.

8. When a scanning laser ophthalmoscope is used to capture images of the fundus, defining medical necessity depends on other factors, as follows:
 - a. If the scanner produces an image of the retina or optic nerve along with other data and imaging for quantitative analysis, it is appropriate to report a single service from the appropriate scanning computerized ophthalmic diagnostic imaging code range (92133-92134).
 - b. If the only required evaluation is a fundus photograph, without the need to quantify the nerve fiber layer and to analyze the data via a computer, it is not considered imaging, even if the photograph was taken with a scanning laser (92250).
9. OCT-angiography may be medically necessary in disorders of the retina and choroid. For these conditions, the procedure is performed independent of other angiography procedures of the eye that involve the use of intravenous dye.

D. DOCUMENTATION

Medical necessity must be supported by adequate and complete documentation in the patient's medical record that describes the procedure and the medical rationale. Documentation requires the following items, all of which must be available upon request. For any retrospective review, a full operative report and/or clinical plan of care is needed.

Every page of the record must be legible and include appropriate patient identification information (e.g., complete name, date(s) of service). Services provided/ordered must be authenticated by the physician. The method used shall be handwritten or electronic signature. Stamped signatures are not acceptable.

Full OCT documentation includes:

1. Physician's order for OCT with medical rationale
2. Date performed
3. Reliability of the OCT (Do not bill an OCT of dubious value)
4. Patient cooperation
5. OCT findings including the interpretation and report. Normal findings may not support the medical necessity of the procedure.
6. Comparison (when applicable) of current and prior OCT impressions
7. Evaluation, diagnosis
8. Impact on treatment, prognosis
9. Do not include copies of the digital images; these must be available upon request.

E. Procedural Detail

CPT Codes	
92132	Scanning computerized ophthalmic diagnostic imaging, anterior segment, with interpretation and report, unilateral or bilateral

92133	Scanning computerized ophthalmic diagnostic imaging, posterior segment, with interpretation and report, unilateral or bilateral, optic nerve
92134	Scanning computerized ophthalmic diagnostic imaging, posterior segment, with interpretation and report, unilateral or bilateral, retina

DISCLAIMER and COPYRIGHTS

This clinical policy is provided for information purposes only and does not constitute medical advice. Versant Health, Inc., and its affiliates (the “Company”) do not provide health care services and cannot guarantee any results or outcomes. Treating doctors are solely responsible for determining what services or treatments to provide to their patients. Patients (members) should always consult their doctor before making any decisions about medical care.

Subject to applicable law, compliance with this clinical policy is not a guarantee of coverage or payment. Coverage is based on the terms of an individual’s particular benefit plan document, which may not cover the service(s) or procedure(s) addressed in this clinical policy. The terms of the individual’s specific benefit plan are always determinative. Every effort has been made to ensure that the information in this clinical policy is accurate and complete, however the Company does not guarantee that there are no errors in this policy or that the display of this file on a website is without error. The company and its employees are not liable for any errors, omissions, or other inaccuracies in the information, product, or processes disclosed herein. Neither the Company nor the employees represent that the use of such information, products, or processes will not infringe on privately owned rights. In no event shall the Company be liable for direct, indirect, special, incidental, or consequential damages arising out of the use of such information, product, or process.

COMPANY’S COPYRIGHT STATEMENT Except for any copyrights described below, this clinical policy is confidential and proprietary, and no part of this clinical policy may be copied, used, or distributed without Versant Health, or its applicable affiliates, expressing prior written approval.

AMA COPYRIGHT STATEMENT CPT© is the 2002-2024 copyright of the American Medical Association. All Rights Reserved. CPT™ is a registered trademark of the American Medical Association. Applicable FARS/DFARS Apply to Government Use. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

RELATED POLICIES AND PROCEDURES	
n/a	

DOCUMENT HISTORY		
Approval Date	Revision	Effective Date
07/10/2017	Initial Policy	07/10/2017
01/23/2018	Annual review; no criteria changes	01/23/2018
12/13/2018	Annual review; no criteria changes	12/13/2018
03/13/2019	Annual review; no criteria changes	03/13/2019
12/8/2019	Name change from SCODI	01/01/2020
10/29/2020	Annual review; addition of criteria for baseline and monitoring of long-term drug therapy	03/01/2020
10/06/2021	Added indication of vigabatrin therapy. Policy opened to all valid, medical ICD-10 codes.	04/01/2022
04/06/2022	Annual review; no criteria changes	05/01/2022
04/12/2023	Annual review; no criteria changes	06/01/2023
04/03/2024	Annual review; no criteria changes	06/01/2024

REFERENCES AND SOURCES

1. Aumann S, Donner S, Fischer J, et al. Optical Coherence Tomography (OCT): Principle and Technical Realization. 2019 Aug 14. In: Bille JF, editor. High Resolution Imaging in Microscopy and Ophthalmology: New Frontiers in Biomedical Optics [Internet]. Cham (CH): Springer; 2019. Chapter 3. PMID: 32091846.
2. Bergeron S, Arthurs B, Sanft DM, et.al. Optical Coherence Tomography of Peri-Ocular Skin Cancers: An Optical Biopsy. *Ocul Oncol Pathol*. 2021 Mar;7(2):149-158. doi: 10.1159/000511188. Epub 2020 Dec 2. PMID: 33981698; PMCID: PMC8077514.
3. Britze J, Frederiksen JL. Optical coherence tomography in multiple sclerosis. *Eye (Lond)*. 2018;32(5):884-888. doi:10.1038/s41433-017-0010-2
4. Daneshmand PG, Rabbani H, Mehridehnavi A. Super-Resolution of Optical Coherence Tomography Images by Scale Mixture Models. *IEEE Trans Image Process*. 2020 Apr 7. doi: 10.1109/TIP.2020.2984896. Epub ahead of print. PMID: 32275595.
5. Do Jeong K, Park JY, Kim BN, et al. Assessment of Choroidal Thickness Inside and Outside of Vascular Arcade in Diabetic Retinopathy Eyes Using Spectral-Domain Optical Coherence Tomography. *Sci Rep*. 2019 Jul 25;9(1):10780. doi: 10.1038/s41598-019-47351-w. PMID: 31346253; PMCID: PMC6658487.
6. Elahi S, Gillmann K, Gasc A, et al. Sensitivity of indocyanine green angiography compared to fluorescein angiography and enhanced depth imaging optical coherence tomography during tapering and fine-tuning of therapy in primary stromal choroiditis: A case series. *J Curr Ophthalmol*. 2019 Jan 17;31(2):180-187. doi: 10.1016/j.joco.2018.12.006. PMID: 31317097; PMCID: PMC6611918.
7. Emre S, Ulusoy MO. Optical coherence tomography angiography findings of the fellow eye of patients with unilateral neovascular age-related macular degeneration OCT-A Evaluation of Fellow Eyes of CNV. *Rom J Ophthalmol*. 2019 Jul-Sep;63(3):231-237. PMID: 31687624;

- PMCID: PMC6820497.
8. Garcia Marin YF, Alonso-Caneiro D, Vincent SJ, et.al. Anterior segment optical coherence tomography (AS-OCT) image analysis methods and applications: A systematic review. *Comput Biol Med.* 2022 Jul; 146:105471. doi: 10.1016/j.compbimed.2022.105471. Epub 2022 Apr 6. PMID: 35533455.
 9. Gumus K, Pflugfelder SC. Anterior Segment Optical Coherence Tomography (AS-OCT) in the Management of Dry Eye. *Int Ophthalmol Clin.* 2017 Spring;57(2):13-22. doi: 10.1097/IIO.000000000000164. PMID: 28282311.
 10. Go MS, Barman NR, Kelly MP, et.al. Overhead Mounted Optical Coherence Tomography in Childhood Glaucoma Evaluation. *J Glaucoma.* 2020 Sep;29(9):742-749. doi: 10.1097/IJG.0000000000001567. PMID: 32496465.
 11. Kolenderska SM, Vanholsbeeck F, Kolenderski P. Fourier domain quantum optical coherence tomography. *Opt Express.* 2020 Sep 28;28(20):29576-29589. doi: 10.1364/OE.399913. PMID: 33114855.
 12. Komma S, Chhablani J, Ali MH, Garudadri CS, Senthil S. Comparison of peripapillary and subfoveal choroidal thickness in normal versus primary open-angle glaucoma (POAG) subjects using spectral domain optical coherence tomography (SD-OCT) and swept source optical coherence tomography (SS-OCT). *BMJ Open Ophthalmol.* 2019 Jul 11;4(1): e000258. doi: 10.1136/bmjophth-2018-000258. PMID: 31414052; PMCID: PMC6668609.
 13. Lee J, Rosen R. Optical Coherence Tomography Angiography in Diabetes. *Curr Diab Rep.* 2016;16(12):123. doi:10.1007/s11892-016-0811-x.
 14. Lisboa R, Leite MT, et.al. Diagnosing pre-perimetric glaucoma with spectral domain optical coherence tomography. *Ophthalmology* 2012; 119:2261-9.
 15. Marmor MF, Keller U, et.al., American Academy of Ophthalmology Revised recommendations on screening for chloroquine and hydroxychloroquine retinopathy, *Ophthalmology* 2011; 118: 415-422.
 16. Masiwa LE, Moodley V. A review of corneal imaging methods for the early diagnosis of pre-clinical Keratoconus. *J Optom.* 2020 Oct-Dec;13(4):269-275. doi: 10.1016/j.optom.2019.11.001. Epub 2020 Jan 6. PMID: 31917136; PMCID: PMC7520528.
 17. Minakaran N, de Carvalho ER, Petzold A, Wong SH. Optical coherence tomography (OCT) in neuro-ophthalmology. *Eye (Lond).* 2021 Jan;35(1):17-32. doi: 10.1038/s41433-020-01288-x. Epub 2020 Nov 25. PMID: 33239763; PMCID: PMC7852683.
 18. Mohammed ISK, Tran S, Toledo-Espielt LA, et.al. The detection of keratoconus using novel metrics derived by anterior segment optical coherence tomography. *Int Ophthalmol.* 2022 Jul;42(7):2117-2126. doi: 10.1007/s10792-021-02210-4. Epub 2022 Jan 6. PMID: 34989951.
 19. O'Bryhim BE, Apte RS, Kung N, et.al. Association of Preclinical Alzheimer Disease with Optical Coherence Tomographic Angiography Findings. *JAMA Ophthalmol.* 2018;136(11):1242–1248. doi:10.1001/jamaophthalmol.2018.3556
 20. Spaide RF, Fujimoto JG, Waheed NK, et al. Optical coherence tomography angiography. *Prog Retin Eye Res.* 2018; 64:1-55. doi: 10.1016/j.preteyeres.2017.11.003.
 21. Veerappan M, El-Hage-Sleiman AM, Tai V, et.al. Age-related Eye Disease Study 2 Ancillary Spectral Domain Optical Coherence Tomography Study Group. Optical Coherence Tomography Reflective Drusen Substructures Predict Progression to Geographic Atrophy in Age-related Macular Degeneration. *Ophthalmology.* 2016 Dec;123(12):2554-2570. doi: 10.1016/j.ophtha.2016.08.047. Epub 2016 Oct 25. PMID: 27793356; PMCID: PMC5125946.
 22. Wang CT, Chang YH, Tan GSW, et.al. Optical Coherence Tomography and Optical Coherence Tomography Angiography in Pediatric Retinal Diseases. *Diagnostics (Basel).* 2023 Apr 18;13(8):1461. doi: 10.3390/diagnostics13081461. PMID: 37189561; PMCID:

PMC10138206.

23. Werner AC, Shen LQ. A Review of OCT Angiography in Glaucoma. *Semin Ophthalmol.* 2019;34(4):279-286. doi: 10.1080/08820538.2019.1620807. Epub 2019 Jun 3. PMID: 31158045.

SOURCES

1. American Academy of Ophthalmology, Drug Induced Maculopathy. “risk of toxicity up to 5 years is under 1%,”. https://eyewiki.org/Drug_Induced_Maculopathy. https://eyewiki.org/Drug_Induced_Maculopathy. Accessed 3/2024.
2. American Academy of Ophthalmology, What Is Optical Coherence Tomography? Apr. 27, 2023. <https://www.aao.org/eye-health/treatments/what-is-optical-coherence-tomography>. Accessed 2/2024.